

**X- RAY POWDER DIFFRACTION DATA FOR
SODIUM MORPHOLYLDITHIOCARBAMATE
TRIHYDRATE $C_5H_8NOS_2Na \cdot 3H_2O$**

^{a)} **Joelis Rodríguez,** ^{b)} **Cristina Diaz,** ^{c)} **Ramón Pomés and** ^{b)} **Roberto Cao**

a) Research Center for Mining and Metallurgy (CIPIMM), Carretera Varona Km 1 ½ Capdevila, Boyeros, Havana Cuba. E-mail: cipimm@chab.minbas.cu

b) Faculty of Chemistry, Havana University, Cuba

c) Albert Schweitzer International University, Geneve, Switzerland

ABSTRACT

Sodium Morpholyldithiocarbamate Trihydrate $C_5H_8NOS_2Na \cdot 3H_2O$ has been investigated by means of X-ray powder diffraction. The title compound is triclinic with unit-cell parameters $a = 6.261 (2)$ $b = 8.897 (3)$ $c = 16.557 (4)$ Å; $\alpha = 118.34 (2)$ $\beta = 99.66 (2)$ $\gamma = 93.32 (3)$ °, $V = 790.2 (3)$ Å³, $Z = 2$, space group P̄1 and $D_x = 1.007 (12)$ g cm⁻³.

KEYWORDS

Crystal characterization, morpholyl dithiocarbamate and X- Ray Powder Diffraction.

INTRODUCTION

Sodium morpholyldithiocarbamate Trihydrate $C_5H_8NOS_2Na \cdot 3H_2O$ (fig. 1) is a compound with several biological properties, such as: antitumoral (1) and antioxidant (2).

This dithiocarbamate anion acts as a S,S type chelating agent and forms stable complexes with representative and transition metals (3, 4) that are generally only slightly soluble or insoluble in water and other polar solvents. The complexes are specially stable with “soft” metals (5).

EXPERIMENTAL

A. Origin of specimen

An ethanol solution of morpholine was added dropwise to an ethanol solution of CS₂ at 0-5 °C (morpholine:CS₂ molar ratio of 1:1). The resulting mixture was treated with Et₂O and an aqueous solution of NaOH for a CS₂:NaOH molar ratio of 1:1. The product was filtered, washed and recrystallized from ethanol; m.p.>300 °C. IR $\nu(\text{cm}^{-1})$: 1460 ($\nu_{\text{C}=\text{N}}$), 981 ($\nu_{\text{C}-\text{S}}$), 542 ($\nu_{\text{C}-\text{S}} + \delta_{\text{SCS}}$). UV: $\lambda_{\text{max}}(\text{nm})$ 263 log ε = 4.18 (CSS $\pi-\pi^*$); 284 log ε = 4.18 (NCS $\pi-\pi^*$), ¹H-NMR (D₂O): δ (ppm) 4.38 (t, 4H, -OCH₂-), J_{H-H}=5.1 Hz); 3.77 (t, 4H, -NCH₂-), J_{H-H}=4.9 Hz).




Fig.1. Scheme of the Sodium Morpholyldithiocarbamate Trihydrate C₅H₈NOS₂Na 3H₂O.

B. Crystal Data

Crystalline powder; color: white; space group: P-1; $a = 6.261(2)$ $b = 8.897(3)$ $c = 16.557(4)$ Å; $\alpha = 118.34(2)$ $\beta = 99.66(2)$ $\gamma = 93.32(3)$ °; $Z = 2$; $D_x = 1.007(12)$ g cm⁻³ and $V = 790.2(3)$ Å³.

C. X-ray powder data

The sample was ground in an agate mortar and the loose powder was pressed into a diffractometer sample holder. The X-ray diffraction pattern was collected using Ni filtered CuK α_1 radiation 1.5406 Å, (K α_2 was eliminated with computer software) on a Philips PW 1710 diffractometer operated at 40 Kv and 30 mA. The alignment of the diffractometer was checked by using a silicon external standard from National Institute of Standards and Technology NIST- SRM- 640b with cell parameter $a = 5.430940(35)$ Å (6) at 25 °C. The powder pattern was recorded at 25 (1) °C from 4 to 70° 2θ using an angular step 0.02° and a counting time of 5 s.

The reported peak heights and positions were extracted by fitting Pearson VII type functions to the diffraction maxima. The positions of all peaks were input in the program for least-square unit cell refinement LSUCRI (7). The starting set of cell parameters for the refinement was taken from the output of the indexing program TREOR90 (8), gave the following cell parameter $a = 6.26$ (1), $b = 8.89$ (2), $c = 16.52$ (2) Å; $\alpha = 118.3$ (2), $\beta = 99.7$ (3), $\gamma = 93.3$ (2)° and figures of merit $M_{20} = 11$ and $F_{20} = 23$ (0.016643, 53) ($\Delta 2\theta$, Npos).

The closed values of the FWHM of 111 peak of the Si standard (FWHM = 0.12° at 28.446°) and that of the 111 peak of the sample (FWHM = 0.14° at 18.976°) indicate a high degree of crystallinity of the specimen. In table I reported the powder diffraction data and in Fig. 2 reported the X-Ray powder pattern of the Sodium Morpholyldithiocarbamate Trihydrate $C_5H_8NOS_2Na \cdot 3H_2O$.

Table 1. Powder diffraction data for Sodium morpholyldithiocarbamate Trihydrate $C_5H_8NOS_2Na \cdot 3H_2O$.

Rad CuK α ($\lambda = 1.5406 \text{ \AA}$) a = 6.261 (2) $\alpha = 118.34$ (2) Z = 2; $D_x = 1.007$ (12) g cm $^{-3}$	Ni filter c = 16.557 (4) \AA $\beta = 99.66$ (2) $\gamma = 93.32$ (3) $^{\circ}$	Sys. Triclinic; Space Group: P $\bar{1}$ V = 790.2 (3) \AA^3 Color White
hkl	2 θ_{obs} ($^{\circ}$)	d $_{\text{obs}}$ (\AA)
0 0 1	6.215	14.208
0 0 2	12.449	7.104
1 0 0, 1 0 -1	14.515	6.098
0 1 1	15.495	5.714
1 -1 0, 1 0 1	17.038	5.200
1 -1 1	17.300	5.128
1 1 -2	18.156	4.882
0 0 3	18.720	4.736
1 -1 -1	18.976	4.679
1 1 0	19.925	4.452
1 0 -3	21.042	4.218
1 0 2	21.242	4.179
0 2 0	23.010	3.862
1 -1 3	23.554	3.774
0 2 -4, 1 -2 2	24.319	3.657
1 0 3	26.199	3.399
0 2 1	26.706	3.335
1 -1 -3	27.214	3.274
1 -1 4, 0 2 -5	28.320	3.148
1 1 -5	28.804	3.097
1 -2 4, 2 0 0	29.387	3.037
0 3 -3, 2 -1 -1	30.209	2.956
2 -1 1, 0 3 -2	30.551	2.924
2 1 -2	30.893	2.892
0 0 5, 2 1 -1	31.455	2.842
1 0 4	31.806	2.811
1 2 1	32.939	2.717
1 3 -4	34.581	2.592
2 -2 2	36.438	2.463
0 3 -6, 2 2 -4	36.681	2.448
1 0 5	37.706	2.384
0 0 6	37.965	2.368
1 -2 6, 0 3 1	38.431	2.340
1 -1 6	39.488	2.280
1 1 -7	39.716	2.268
2 1 -6, 2 2 0	40.400	2.231
1 -4 2	42.983	2.103
1 -2 7, 2 -2 5	43.720	2.069
0 0 7	44.602	2.030

1 -2 -5	47.036	1.9304	9
3 -2 1, 2 -3 -2	47.400	1.9164	8
2 0 5, 3 -2 2	48.415	1.8786	5
3 -2 -2, 2 2 -8 , 2 1 4, 3 2 .	49.328	1.8460	4
0 4 8, 2 -3 6, 2 -1 6,	49.756	1.8310	10
2 -3 -3, 1 1 6, 3 1 -6	50.996	1.7894	3
0 0 8, 2 -4 5	51.408	1.7760	4
2 3 -8, 0 5 -5	51.519	1.7724	4
3 -3 1, 1 1 -9	52.138	1.7529	3
3 -1 -6, 3 3 -6, 3 -2 5	56.582	1.6253	5
1 0 -9	56.882	1.6174	4
1 -3 -5, 3 3 -2, 1 1 7 , 2 -4	57.505	1.6014	3
0 0 9, 2 -3 8	58.395	1.5788	3
4 0 -1, 4 -1 -1, 2 4 -9	59.475	1.5530	5
3 4 4, 4 0 -4	60.394	1.5315	3
2 3 3	62.579	1.4832	4
1 -1 -9, 2 5 -3, 0 4 -11	63.923	1.4552	3
3 -1 -8	66.287	1.4089	3




Fig.2. X-ray diffraction pattern of the Sodium Morpholyldithiocarbamate Trihydrate $C_5H_8NOS_2Na \cdot 3H_2O$.

RESUMEN

El compuesto Morfolil Ditiocarbamato trihidratado $C_5H_8NOS_2Na \cdot 3H_2O$ fue estudiado a través de difracción de rayos- X, por el método de polvo. El mismo cristaliza en el sistema triclínico, en el grupo espacial $P\bar{1}$, con parámetros de celda: $a = 6.261 (2)$ $b = 8.897 (3)$ $c = 16.557 (4) \text{ \AA}$; $\alpha = 118.34 (2)$ $\beta = 99.66 (2)$ $\gamma = 93.32 (3)^\circ$, $V = 790.2 (3) \text{ \AA}^3$, $Z = 2$ y $D_x = 1.007 (12) \text{ g cm}^{-3}$.

PALABRAS CLAVES

Caracterización cristalográfica, Morfolil ditiocarbamate, Difracción de Rayos- X por el método de polvo.

BIBLIOGRAPHY

Mainardi, V., J. L. Bello, J. M. Sanfiz & F. Torrez. Rev. Cubana Oncol. 1990, 6, 107-111.

Cao, R., A. Fragoso & R. Villalonga. Monatsh. Chem. 1996, 127, 775-782.

Beyer, W. & R. D. Ott. Mikrochim. Acta 1965, 1130-1135.

Aravamudan, G., D. H. Brown & D. Venkappayya. Journal Chem. Soc. 1971, (A) 2744-2747.

Sastri, V. S., K. I. Aspila & C. L. Chakrabarti. Canadian Journal of Chemistry. 1969, 47 2320-2323.

Pomés, R.: "Análisis de fases cuantitativo usando difracción de Rayos - X", Editorial Quetzatcoal, Mex. 1997, 140.

Garvey, R.: "LSUCRI PCUnit Cell Refinement with Indexing for the Personal Computer", Powder Diffraction (1), 1986, 114-116.

Werner, P. E., L. Eriksson & M. Westdahl: "TREOR, a semi exhaustive trial-and-error powder indexing program for all symmetries", Journal of Applied Crystallography, (18), 1985, 367-370.

Recibido marzo del 2002, aceptado abril del 2002.